

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 303/38, 311/48	A1	(11) International Publication Number: WO 97/23448 (43) International Publication Date: 3 July 1997 (03.07.97)
(21) International Application Number: PCT/US96/19532 (22) International Filing Date: 9 December 1996 (09.12.96) (30) Priority Data: 08/577,425 22 December 1995 (22.12.95) US (71) Applicant: MINNESOTA MINING AND MANUFACTURING COMPANY [US/US]; 3M Center, P.O. Box 33427, Saint Paul, MN 55133-3427 (US). (72) Inventors: HOWELLS, Richard, D.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). LAMANNA, William, M.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). FANTA, Alan, D.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). WADDELL, Jennifer, E.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). (74) Agents: SCHULTE, Daniel, C. et al.; Minnesota Mining and Manufacturing Company, Office of Intellectual Property Counsel, P.O. Box 33427, Saint Paul, MN 55133-3427 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PREPARATION OF BIS(FUOROALKYLENESULFONYL)IMIDES AND (FUOROALKYLSULFONYL) (FLUOROSULFONYL)IMIDES (57) Abstract A method for preparing fluoroalkylsulfonyl imides by reacting a fluoroalkylsulfonamide with a fluoroalkylsulfonyl halide or a fluorosulfonyl halide in the presence of a non-nucleophilic base. One reaction is: $Z-R_f-SO_2NH_2 + Z-R'_f-SO_2-X + 2B \Rightarrow (Z-R_f-SO_2)(Z-R'_f-SO_2)N^+BH^+ + BHX$, where each Z is a fluorine atom or a polymerizable organic functional group, R_f and R'_f are fluoroalkylene groups optionally containing catenary oxygen or nitrogen, X is a halogen atom, and B is a non-nucleophilic base. Unsymmetrical imides and polymeric imides can be prepared.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Preparation of Bis(fluoroalkylenesulfonyl) Imides and
(Fluoroalkylsulfonyl) (Fluorosulfonyl) Imides

Technical Field

This invention relates to a method for
5 preparing bis(fluoroalkylsulfonyl)imides,
(fluoroalkylsulfonyl) (fluorosulfonyl)imides and
derivatives thereof.

Background

10 Bis(Fluoroalkylsulfonyl)imides,
(fluoroalkylsulfonyl) (fluorosulfonyl)imides and
derivatives thereof are known to be useful in many
chemical processes and formulations. In solution, the
imides behave as strong acids due to the presence of
15 the electron-withdrawing fluoroalkylsulfonyl or
fluorosulfonyl groups and exhibit exceptional stability
under a variety of conditions. These compounds have
been found to be useful, for example, as fuel cell
electrolytes, as esterification catalysts and as
20 polymerization catalysts for cationically-sensitive
monomers such as epoxy resins.

The salts of bis(fluoroalkylsulfonyl)imides
and (fluoroalkylsulfonyl) (fluorosulfonyl)imides have
been found useful as electrolytes in advanced high
25 voltage batteries. In particular, lithium
bis(trifluoromethylsulfonyl)imide $(\text{CF}_3\text{SO}_2)_2\text{N}^-\text{Li}^+$, shows
excellent conductivity and stability, but can be
corrosive toward aluminum. Certain higher homologues
of this compound show comparable performance as battery
30 electrolytes, and have decreased the corrosion toward
aluminum.

There are several known methods for preparing
some classes of bis(fluoroalkylsulfonyl)imides.
Bis(perfluoroalkylsulfonyl)imides can be prepared by

the reaction of an alkali metal salt of an N-trimethylsilyl perfluoroalkylsulfonamide with a perfluoroalkylsulfonyl fluoride in a polar organic solvent. This route requires several synthetic steps, with the isolation of intermediate compounds, and the overall yields are often low. A second method involves the reaction of an alkali metal salts of a perfluoroalkylsulfonamide with perfluoroalkylsulfonyl fluoride in a polar organic solvent. Here too, the yields are low. A third method involves reacting a perfluorinated sulfonic anhydride with urea and a sulfonic acid. The products of this reaction are then dissolved in water and the addition of tetrabutylammonium bromide precipitates the tetrabutylammonium imide. This method is not satisfactory for the large scale production of imides because the overall yields are low and the sulfonic anhydride precursors are not easily accessible. A fourth method of making metal salts of sulfonimides involves the reaction of a metal nitride with perfluoroalkyl sulfonyl halide. However this method cannot be used to make unsymmetrical imides and requires handling metal nitrides which are sensitive to air.

Thus there remains a need for an efficient method of preparing bis(fluoroalkylsulfonyl)imides, (fluoroalkylsulfonyl)(fluorosulfonyl)imides, and derivatives thereof.

Disclosure of the Invention

The present invention provides an efficient, high yield method of preparing bis(fluoroalkylsulfonyl)imides, (fluoroalkylsulfonyl)(fluorosulfonyl)imides as well as derivatives such as the organic or inorganic salts, and


N-halo, N-alkyl or N-silyl compounds; hereinafter described as "imides" for the sake of brevity. The method uses readily available starting materials and allows the preparation of the imides in a single step, without the isolation and purification of intermediate compounds. The method of this invention also features the preparation of unsymmetrical imides and polymers having imide groups in the polymer chain.

By this method, a fluoroalkylsulfonamide is contacted with a fluoroalkylsulfonyl halide or a fluorosulfonyl halide, in the presence of a sufficient amount of a non-nucleophilic base at a temperature and for a time sufficient to yield the imide salt. Preferably, the fluoroalkylsulfonamide is prepared *in situ* by contacting a fluoroalkylsulfonyl halide with anhydrous ammonia.

Detailed Description

In one aspect, the method of the invention may be represented by the following Scheme 1:


Scheme 1



wherein R_f and R'_f are fluoroalkylene groups having from 1 to 12 carbon atoms, optionally containing one or more catenary oxygen or nitrogen atoms, each Z is a fluorine atom or a polymerizable organic functional group, X is a halogen atom and B is a non-nucleophilic base. BH' represents an cation derived from the non-nucleophilic base B, e.g. an ammonium cation, but other cations may be used. In the case of symmetrical imides, the sulfonamide is preferably generated *in situ* by the reaction of a sulfonyl halide with anhydrous ammonia. The sulfonamide then reacts with another


molecule of sulfonyl halide to produce the bis(fluoroalkylenesulfonyl)imide as shown in Scheme 2.

Scheme 2




when sulfonamides are used as the starting compounds in the preparation of bis(fluoroalkylenesulfonyl)imides according to the method of this invention, either the sulfonamide *per se*, or a salt of the sulfonamide may be used. As used herein, the term "sulfonamide" represents both the sulfonamides and the salts thereof. Thus, the use of fluoroalkylenesulfonamide salts may be represented in Scheme 3.

Scheme 3




In another aspect of the invention, polymeric bis(fluoroalkylenesulfonyl)imides may be prepared by contacting equimolar amounts of difunctional fluoroalkylene sulfonyl halides and fluoroalkylene sulfonamides in the presence of a non-nucleophilic base. The difunctional sulfonyl halides or sulfonamides may be fluoroalkylene disulfonyl halides or fluoroalkylene disulfonamides. Alternatively, a difunctional sulfonamidofluoroalkylenesulfonyl halide may be used. The preparation of polymeric imides is shown in Schemes 4-6.

Scheme 4




Scheme 5



5

Scheme 6




In Scheme 6, the sulfonamide is generated *in situ* by the reaction between a sulfonyl halide and ammonia in the presence of a non-nucleophilic base B. The sulfonamide thus produced reacts with a sulfonyl halide from another molecule of $\text{X-SO}_2\text{-R}_f\text{-SO}_2\text{-X}$ to produce the imide. Where sulfonamides are used as starting compounds, either the sulfonamide per se can be used, or a salt as previously described. Preferably $\text{X-SO}_2\text{-R}_f\text{-SO}_2\text{-X}$ is chosen so that the R_f group is sufficiently large to make intramolecular cyclization unfavorable. Cyclization would be unfavorable when the resulting cyclic compound would have a ring size \leq four ring atoms or \geq nine ring atoms.

In the preparation of polymeric imides according to Schemes 4-6 it is preferable that the reagents be of the highest possible purity (e.g., $> 99\%$ by weight) and have a high level of difunctionality (e.g., $> 99\%$) so that the reaction proceeds to high levels of conversion (e.g., $> 98\%$) with minimal side reactions in order to maximize the molecular weight of the final polymer.


(Fluoroalkylsulfonyl)(fluorosulfonyl)imides can be prepared by the method of this invention using a fluoroalkylsulfonamide and a fluorosulfonyl halide according to Scheme 7.

Scheme 7



The fluoroalkylsulfonamide is preferably
 5 generated *in situ* by the reaction between a
 fluoroalkylsulfonyl halide and ammonia, and
 subsequently reacted with the fluorosulfonyl halide to
 produce the imides. As previously described, the
 fluoroalkylsulfonamide per se may be used, or a salt as
 10 shown in Scheme 8.


Scheme 8




15

Difunctional fluoroalkylenesulfonamides may
 also be reacted with monofunctional
 fluoroalkylenesulfonyl halides, or conversely,
 difunctional fluoroalkylenesulfonyl halides may be
 20 reacted with monofunctional fluoroalkylenesulfonamides
 to produce dimeric bis(fluoroalkylenesulfonyl)imides,
 as shown in Schemes 9 and 10.

Scheme 9



Scheme 10



The fluoroalkylene groups, R_f and R'_f , of the
 fluoroalkylenesulfonyl halides and the
 fluoroalkylenesulfonamides may be linear, branched,
 35 cyclic or acyclic, or combinations thereof.

Additionally one or more of the fluorine or hydrogen atoms of the fluoroalkylene group may be replaced by chlorine or bromine atoms. In the above formulae shown in Schemes 1 to 10, one or more non-adjacent carbon atoms of the fluoroalkylene groups R_f and R'_f may be replaced by catenary divalent oxygen or trivalent nitrogen atoms.

When the fluoroalkylene group is acyclic it has the general formula $C_mH_{(2m-a)}F_a$, where m is 1 to 12 and a is at least one. When the fluoroalkylene group contains a cyclic group it has the general formula $C_mH_{(2m-2-a)}F_a$, where m is 1 to 12 and a is at least one. Preferably the sum of carbon atoms in the two fluoroalkylene groups of the bis(fluoroalkylenesulfonyl)imide is 3 to 12 carbon atoms

Preferably, the fluoroalkylene group is substantially fluorinated so that the number of fluorine atoms is greater than the number of hydrogen atoms present. Most preferably the fluoroalkylene group is a linear or branched, cyclic or acyclic perfluoroalkylene group containing 1 to 10 carbon atoms and the sum of carbon atoms, in the two fluoroalkylene groups of the bis(fluoroalkylenesulfonyl)imide is 3 to 12 carbon atoms.

The halide X may be F , Br or I , but is preferably F . When Z is a polymerizable organic functional group it allows the preparation of polymers having pendant bis(fluoroalkylenesulfonyl)imide moieties on the polymer chain.

Fluoroalkylenesulfonylfluorides having polymerizable functional groups have been described by Gard et al., J. Fluorine Chem. 66, 105(1994), Gard et al., Coordination Chemistry Reviews 112, 47(1992), Gard et al., J. Fluorine Chem. 49, 331(1990), Gard et al. J.

- Fluorine Chem. 43, 329(1989), Gard et al., J. Fluorine Chem. 67, 27(1994), Gard et al. J. Fluorine Chem. 55, 313(1991), Gard et al. J. Fluorine Chem. 38, 3(1988), Gard et al., Inorg. Chem., 29, 4588(1990) U.S. 5,414,117 (Armand) and U.S. 5,463,005 (Desmarteau). Polymers prepared from fluoroalkylenesulfonylfluorides having polymerizable functional groups have been described in DesMarteau, Novel Fluorinated Acids for Phosphoric Acid Fuel Cells, Gas Research Institute Report #GRI-92/0385, July 1992, and J. Fluorine Chem. 72, 203 (1995).

- Z may be any organic functional group that may undergo grafting or polymerization by an addition or step-growth mechanism, including cationic, anionic, free-radical and Z-N polymerization or polycondensation. Z may be chosen from those groups containing double bonds, for example vinyl, allyl, vinyl-benzyl or acryloyl groups. Z may also be chosen from those groups containing oxirane, oxetane, azetidine, or aziridine functional groups. Z may also be chosen from those groups containing alcohol, amine, isocyanate, or trialkoxysilyl functional groups.

- When Z contains functional groups whose reactivity could interfere with reactions for preparing the imides, Z can be protected by reactants that are reversibly bound to it. For example, a double bond may be protected as a di-halo derivative and subsequently dehalogenated. Examples of Z include $\text{CH}_2=\text{CH}-$, $\text{CH}_2=\text{CHO}-$, $\text{CH}_2=\text{CHCH}_2\text{O}-$, $\text{CH}_2=\text{CHCO}-$, $(\text{CH}_2=\text{CHCH}_2)_2\text{NCO}-$ (prepared as described in Armand et.al, US 5,414,117), $(\text{CH}_2=\text{CHCH}_2)_2\text{NSO}_2-$, $\text{CF}_3=\text{CFO}-$, $\text{CF}_2=\text{CF}-$, $\text{CH}_2=\text{CHCO}_2\text{CH}_2\text{CH}_2-$, $-\text{O}-\text{C}_6\text{H}_4-\text{CH}=\text{CH}_2$, $-\text{C}(\text{O})\text{OCH}_2\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{C}(\text{O})\text{NCH}_2\text{CH}_2\text{OC}(\text{O})\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{SO}_2\text{O}-\text{C}_6\text{H}_4-\text{CH}=\text{CH}_2$, $-\text{OCH}_2\text{CHOH}-\text{CH}_2\text{OH}$, $-\text{C}(\text{O})\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{C}(\text{O})\text{OCH}_2\text{CH}=\text{CH}_2$, and $-\text{OCH}_2\text{C}\equiv\text{CH}$.

Perfluoroalkylsulfonyl fluorides and perfluoroalkylenedisulfonyl fluorides used as precursors to the imide salts of this invention can be prepared by a variety of methods known in the art as described, for example, in U.S. Patents 3,542,864; 5,318,674; 3,423,299; 3,476,753, 3,776,960, 2,519,983, 3,951,762; 3,623,963; 2,732,398 S. Temple, J.Org. Chem., 33(1), 344 (1968), D.D. DesMarteau, Inorg. Chem., 32, 5007 (1993), and Chemistry of Organic Fluorine Compounds, M. Hudlicky, ed., 2nd ed. PTR Prentice Hall (New York), pp. 73-6.

Perfluoroalkylsulfonyl fluorides are available by the electrochemical fluorination of the corresponding alkylsulfonyl fluorides as described in U.S. 2,732,398 (Brice et al.). See also T. Abe et al., in Preparation, Properties and Industrial Applications of Organofluorine Compounds, R.E. Banks(Ed), pages 37-9, John Wiley & Sons, New York (1982). Routes to perfluoroalkanesulfonyl chlorides are described by P.J. Stag et al. in Synthesis 1982, 85 and by R.N. Haszeldine et al., in J. Chem. Soc. 1955, 2901. The chlorides of ω -hydrofluorinated sulfonic acids are described by Coffman and Raasch in J. Org. Chem., 14, 747 (1949), by Weiyuan et al. in Chemistry 2, 31 (1987), and by Huang, J. Fluorine Chem. 32, 179 (1986).

Ammonia, when used in the method of this invention, should be anhydrous to prevent side reactions from occurring. To avoid the hazards of handling liquid ammonia, it is often convenient to generated anhydrous ammonia *in situ* by reacting ammonium salts of the formula NH_4^+X^- with one equivalent of a non-nucleophilic base. Here X may be any halide, or any other non-reactive anion.

For purposes of this description, the term "non-nucleophilic base" B means a base which does not

undergo an irreversible reaction with the sulfonyl halide group. This reaction, when it occurs will reduce the yields of the desired imide. The non-nucleophilic base may be an organic or inorganic base, but is preferably an organic aprotic base. Examples of suitable organic non-nucleophilic bases include alkylamines, for example triethyl amine, trimethyl amine, tripropyl amine and diisopropylethyl amine, pyridines, alkyl pyridines and dialkylaminopyridines, alkyl piperadines, dialkyl piperazines, N-alkyl pyrrolidines, N-alkylpyrroles N-alkylimidazoles, amidines, for example 1,5-diazabicyclo[4.3.0]non-5-ene (DBN) or 1,8-diazabicyclo[5.4.0]dec-7-ene (DBU), or guanidines.









It is preferable to prepare the bis(fluoroalkylenesulfonyl)imides neat, i.e. in the absence of any solvent. Generally a solvent is not necessary, the reactants are sufficiently soluble in each other to produce the imides. In some cases, solvents may be added to enhance the solubility of the reactants, such as when preparing polymeric imides from difunctional sulfonyl halides and/or sulfonamides. Any solvent that is non-reactive toward the starting sulfonyl halides and sulfonamides may be used, but polar aprotic solvents are preferred. Any solvent used should be anhydrous, (e.g., less than 1% water) to avoid competing hydrolysis of the sulfonyl halides. Examples of suitable aprotic liquids include linear ethers such as diethyl ether, diethylene glycol dimethyl ether, and 1,2-dimethoxyethane; cyclic ethers such as tetrahydrofuran, 2-methyltetrahydrofuran, dioxane, dioxolane, and 4-methyldioxolane; nitriles such as acetonitrile and benzonitrile; nitro compounds such as nitromethane or nitrobenzene; amides such as N,N-dimethylformamide, N,N-diethylformamide, and N-







methylypyrrolidinone; sulfoxides such as dimethyl
sulfoxide; sulfones such as dimethylsulfone,
tetramethylene sulfone, and other sulfolanes;
oxazolidinones such as N-methyl-2-oxazolidinone and
5 mixtures thereof.

The method of this invention can be carried
out by introducing the reactants into a vessel, which
can be made of glass or other corrosion-resistant
material. Preferably the vessel is suitable for use at
10 elevated pressures caused by the high vapor pressure(s)
of certain reagents (e.g., NH_3). The method of this
invention can generally be carried out by heating the
contents of the vessel at temperatures from about -25°C
to about 150°C , preferably from 25°C to about 100°C and
15 at a pressure equal to or greater than atmospheric.
Generally pressures in excess of those generated by the
vapor pressures of the reagents at the reaction
temperatures is not required.

When preparing non-polymeric imides from
20 fluoroalkylsulfonamides generally at least one molar
equivalent of the sulfonyl halide is used. Preferably
a slight stoichiometric excess of the sulfonyl halide
is used. When preparing imides from ammonia, as shown
in Scheme 2, a stoichiometric excess of sulfonyl
25 halide, relative to the amount of ammonia, is generally
preferred. The amount of non-nucleophilic base used is
generally in excess of the stoichiometric amount shown
in Schemes 1 through 6. Preferably the amount of base
is less than a 100% excess of the stoichiometric amount
30 shown in the Schemes. Most preferably amount of base
is less than a 50% excess of the stoichiometric amount
shown in the Schemes to maximize the volume efficiency
of the reaction.

Representative imides that can be prepared by
35 the method of this invention include:





The imides shown (without a corresponding cation for illustrative purposes) can be used to prepare derivatives such as alkali, alkaline earth and other metal salts, N-silyl imides, N-alkyl and N-fluoro imides. For example, conjugate acids of the imides may be treated with an inorganic base having the desired cation M', such as NaOH, LiOH, KOH, NH₄OH, Ag₂CO₃, or CuO, to produce the corresponding N-Na, N-Li, N-K, N-NH₄, N-Ag or N-Cu salts. The imides may be contacted with fluorine under direct fluorination conditions to produce the N-F derivatives. The silver imide salts may be contacted with a trialkylsilyl chloride, trialkylstannyl chloride, or an alkyl halide to produce the N-trialkylsilyl, N-trialkylstannyl or N-alkyl derivatives respectively. Other derivatives can be prepared by means known to those skilled in the art.

The invention is further illustrated by the following Examples. In the examples, the term "reduced pressure" means the vacuum obtained by means of a commercially available laboratory distillation apparatus using an aspirator connected to a water pipe or other fluid stream for aspiration, such as a rotary evaporator apparatus.

Example 1

Lithium(trifluoromethanesulfonyl)

(perfluorobutanesulfonyl)imide

Under a nitrogen atmosphere, a dry, 500 mL Fisher-Porter pressure bottle equipped with a stainless steel pressure head and magnetic stirrer was charged with 35.00 g $\text{CF}_3\text{SO}_2\text{NH}_2$ (prepared, e.g., as described in Foropoulos and DesMarteau, *Inorg. Chem.*, 23:3720-23 (1984)), 98 mL anhydrous triethylamine and 74.55 g $\text{C}_4\text{F}_9\text{SO}_2\text{F}$. The pressure vessel was sealed and the reaction mixture heated to 90°C with stirring for 17 hours. The volatile components of the reaction mixture were evaporated under reduced pressure at 70°C then the reaction mixture was treated with a mixture of 700 mL water and 700 mL methylene chloride with stirring to form a two phase mixture. The methylene chloride phase was then separated, washed with two 700 mL portions of water, dried over anhydrous MgSO_4 , filtered and then evaporated under reduced pressure to yield a dark red liquid. This liquid was combined with 500 g of polyphosphoric acid (Aldrich Chemical Co.) in a short path vacuum distillation apparatus equipped with an ambient air-cooled condenser, and then vacuum distilled at ca. 0.1 Torr. The distillate (96.5 g, corresponding to $\text{HN}(\text{SO}_2\text{CF}_3)(\text{SO}_2\text{C}_4\text{F}_9)$, which solidified in receiver at 0°C, was further purified by sublimation at 70°C, 1.0 Torr and then added in portions to a slurry of 52 g of Li_2CO_3 (99.997% purity, from Aldrich Chemical Co.) in 800 mL of methyl t-butyl ether with stirring at room temperature. After about 1 hour CO_2 evolution subsided and the reaction mixture was heated to about 40°C for 3.5 hours to complete the neutralization. The mixture was filtered by gravity through filter paper and then filtered again by suction through a 0.22 micron Tefsep® Teflon membrane (Micron Separations Inc., Westboro, Massachusetts) to remove particulates. The filtrate

was evaporated under reduced pressure at 25-80°C to yield a clear, colorless oil. Two 250 mL portions of toluene were combined with the oil and the mixture evaporated at 40-70°C, 20 Torr after each toluene addition, causing the oil to eventually solidify giving a fine white powder. The solid was transferred to a glass jar and dried in vacuo at 100°C, 10^{-2} Torr for 24 hours to yield 77.0 g of the anhydrous title salt. The structure of the product was confirmed by ^1H and ^{19}F NMR spectroscopy which indicated that the purity of the salt was greater than 99% by weight.

Example 2

Lithium (trifluoromethanesulfonyl)
(perfluoroethanesulfonyl)imide

Under a nitrogen atmosphere, a dry 3L flask equipped with a dry ice condenser, mechanical stirrer, and a sparging tube was charged with 500 mL anhydrous methyl t-butyl ether (MTBE) and cooled below 0°C in a dry ice bath. To the cold solvent was added 500 g of crude $\text{CF}_3\text{CF}_2\text{SO}_2\text{F}$ mixture (containing ca. 23% $\text{CF}_3\text{CF}_2\text{SO}_2\text{F}$ by weight in C6-C8 perfluoroalkane solvent). Gaseous anhydrous ammonia (194 g) was gradually introduced through the sparging tube into the resulting mixture at ca. 0°C with vigorous stirring. Once ammonia addition was complete, the reaction was allowed to proceed for an additional hour after which the dry ice bath was removed and the reaction solution gradually warmed to room temperature with stirring. Excess ammonia was allowed to evaporate while stirring overnight at room temperature. The reaction solution was cooled again to 0°C and treated with 83.9 g $\text{LiOH}\cdot\text{H}_2\text{O}$ dissolved in 750 mL of water with stirring. The mixture was filtered through filter paper by gravity to remove LiF

precipitate. To the filtrate was added concentrated aqueous HCl in portions with stirring at 0°C until the pH of the mixture was 0.

The resulting two-phase mixture was separated
5 into separate aqueous and ether phases. The aqueous phase was extracted with two additional 400 mL portions of MTBE. The ether fractions were combined, extracted with two 500 mL portions of water, and then dried over anhydrous MgSO_4 overnight. The ether solution was
10 filtered by gravity through filter paper and then by suction through a 0.22 micron Tefsep™ membrane (Micron Separations Inc.). The filtrate was concentrated by rotary evaporation, hexane was added, and the solution concentrated again. This was repeated until the
15 product $\text{CF}_3\text{CF}_2\text{SO}_2\text{NH}_2$ crystallized from solution as a white solid. The product (Crop 1) was isolated by suction filtration through a glass frit giving 33.0 g. A second crop of crystals (20.2 g) was recovered in the same manner from the filtrate by concentrating further
20 followed by filtration. The remaining filtrate was evaporated to dryness at 40°C, 20 torr to yield a third crop of relatively crude product (26.8 g). All product fractions (Crops 1-3) were dried at 10^{-2} torr at room temperature for ca. 15 min to give a combined yield of
25 80.0 g. Quantitative ^1H and ^{13}C NMR analysis of the product from Crop 2 indicated that it was the desired product with a purity of 98% by weight.

Under a nitrogen atmosphere, a dry Parr 4560-Series Benchtop Mini Reactor equipped with a 600 mL
30 capacity Monel reactor cylinder, mechanical stirrer, thermocouple and heating mantle was charged with 53 g $\text{CF}_3\text{CF}_2\text{SO}_2\text{NH}_2$ (Crops 1 + 2) and 152 mL anhydrous triethylamine. While cooling the reactor at -78°C in dry ice, 55.2 g of gaseous $\text{CF}_3\text{SO}_2\text{F}$ (ca. 94% purity) was
35 condensed into the reactor with stirring. The reactor

was sealed and the temperature of the reaction mixture gradually raised to 90°C with vigorous stirring, then held at 90°C with stirring for a total of 24 hours. The volatile components of the reaction mixture were

5 evaporated under reduced pressure at 70°C, then the residue was treated with a mixture of 700 mL water and 700 mL methylene chloride with stirring to form a two phase mixture. The methylene chloride phase was then separated, washed with two 700 mL portions of water,

10 dried over anhydrous MgSO_4 , filtered and then evaporated under reduced pressure to yield a dark red liquid. This liquid was combined with 600 g of polyphosphoric acid (Aldrich Chemical Co.) in a short path vacuum distillation apparatus equipped with an ambient air-

15 cooled condenser, and then vacuum distilled at ca. 15 torr, 85-88°C. The distillate (78.1 g, corresponding to $\text{HN}(\text{SO}_2\text{CF}_3)(\text{SO}_2\text{C}_2\text{F}_5)$), which solidified in receiver at 0°C, was converted to the corresponding lithium salt, and further purified, using essentially the procedure

20 described in Example 1 to yield 75.0 g of the anhydrous title salt. The structure of the product was confirmed by ^1H and ^{19}F NMR spectroscopy which indicated that the purity of the salt was 98% by weight.

Example 3

Lithium Bis(perfluoroethanesulfonyl)imide

5 As in Example 2, a dry Parr 4560-Series
Benchtop Mini Reactor was charged with 155 mL anhydrous
triethylamine. While cooling the reactor at -78°C in
dry ice, gaseous $\text{C}_2\text{F}_5\text{SO}_2\text{F}$ (100.0 g, > 99% purity)
followed by anhydrous ammonia (3.51 g) were condensed
10 into the reactor with stirring. The reactor was sealed
and the temperature of the reaction mixture gradually
raised to 90°C with vigorous stirring. The reaction
mixture was held at 90°C with stirring for a total of
20.5 hours. The intermediate product, $\text{HN}(\text{SO}_2\text{C}_2\text{F}_5)_2$, was
15 isolated as previously described, then added to a
slurry of Li_2CO_3 (45.7 g, 99.997% purity) in 800 mL of
methyl t-butyl ether to produce the lithium salt. This
lithium salt was isolated using essentially the
procedure described in Example 1 to yield 73.34 g (92%
20 yield based upon NH_3) of the anhydrous title salt. The
structure of the product was confirmed by ^1H and ^{19}F NMR
spectroscopy which indicated that the purity of the
salt was 99.9% by weight.

25

Example 4

Lithium Bis(perfluorobutanesulfonyl)imide

As in Example 1, a dry, 500 mL Fisher-Porter
pressure bottle was charged with anhydrous
30 triethylamine (100.5 g) and $\text{C}_4\text{F}_9\text{SO}_2\text{F}$ (145.0 g). The
pressure vessel was sealed, cooled to -78°C in dry ice
and then charged with of anhydrous ammonia (3.0 g) with
stirring. After warming gradually to room temperature,
the reaction mixture was heated to 90°C with stirring
35 for 24 hours. The intermediate product, $\text{HN}(\text{SO}_2\text{C}_4\text{F}_9)_2$,

was isolated as previously described, then added to a slurry of Li_2CO_3 (40.0 g, 99.997% purity) in 800 mL of methyl t-butyl ether to produce the lithium salt. This lithium salt was isolated using essentially the
5 procedure described in Example 1 to yield 90.8 g (86% yield based upon NH_3) of the anhydrous title salt. The structure of the product was confirmed by ^1H and ^{19}F NMR spectroscopy which indicated that the purity of the salt was 98% by weight.

10

Example 5

Lithium Bis(perfluoropropanesulfonyl)imide

As in Example 4, the title compound was
15 prepared using anhydrous triethylamine (100.5 g), $\text{C}_3\text{F}_7\text{SO}_2\text{F}$ (100.0 g, 68:32 iso:normal isomer ratio), and anhydrous ammonia (3.0 g). A total of 40.8 g of the anhydrous title salt was recovered as a light pink solid. Quantitative analysis by ^1H and ^{19}F NMR
20 spectroscopy indicated that the product contained the following major components in order of decreasing weight percent: $\text{Li}^+\text{N}(\text{SO}_2\text{C}_3\text{F}_7)_2$, 85.1%, 39:61 iso:normal C_3F_7 ratio; $\text{Li}^+\text{N}(\text{SO}_2\text{i-C}_3\text{F}_7)(\text{SO}_2\text{F})$, 8.7%; $\text{Li}^+\text{N}(\text{SO}_2\text{n-C}_3\text{F}_7)(\text{SO}_2\text{F})$, 4.2%.

25

Example 6

Synthesis of Lithium trifluoromethylsulfonyl
perfluoro(dimethylaminoethylsulfonyl) imide
30 $\text{Li}^+\text{N}(\text{SO}_2\text{CF}_3)(\text{SO}_2\text{C}_2\text{F}_4\text{N}(\text{CF}_3)_2)$

The hydrocarbon precursor 2-dimethylaminoethylsulfonylfluoride was made by the addition of dimethylamine to vinylsulfonyl fluoride as
35 described in Krutak et al., J. Org. Chem., 44(2):3847-58 (1979). The hydrocarbon was perfluorinated by electrochemical fluorination. An electrochemical cell

of the type described in U.S. Patent No. 2,519,983 (Simons) was filled with 1500 cm³ anhydrous HF. The cell was operated at 31 psig (214 Kpa) and at an average temperature of 56°C. During the course of the 129 hour run, 456 grams of fluorinated product was produced from 650 grams (4.2 mole) of starting material. Analysis of the fluorocarbons was done using gas chromatography/FTIR to confirm structures and yields. The resulting fluorochemical was distilled in a 3-plate distillation column to a main cut of 374 grams (1.1 moles, 27% molar yield) corresponding to perfluorodimethylaminoethylsulfonyl fluoride ((CF₃)₂NC₂F₄SO₂F).

To a dry flask equipped with a reflux condenser and magnetic stirrer was charged 1.13 g CF₃SO₂NH₂ (prepared, e.g., as described in Foropoulos and DesMarceau, Inorg. Chem., 23:3720-23 (1984)), 20 ml. anhydrous triethylamine (distilled from LiAlH₄) and 2.67g (CF₃)₂NC₂F₄SO₂F. The reaction mixture was heated to 70°C under a nitrogen atmosphere with stirring for 17 hours, after which volatile components were removed under reduced pressure. The residue was treated with a mixture of 40 mL water and 40 mL methylene chloride with stirring to form a two phase mixture. The methylene chloride phase was then separated, washed with two 40 mL portions of fresh water to remove water-soluble components, dried over anhydrous MgSO₄, filtered and then evaporated under reduced pressure to yield a dark red liquid. The liquid residue was combined with 20 mL of concentrated (at least 98 weight percent H₂SO₄) sulfuric acid in a short path distillation apparatus equipped with a dry-ice cooled condenser, and then vacuum distilled at 10⁻³ torr, 60°C. The distillate was dissolved in 50 mL of diethyl ether, after which the resulting ether solution was treated with excess

lithium carbonate for 2 hours with stirring at room temperature, filtered and the filtrate evaporated under reduced pressure to yield a clear, colorless oil.

Approximately 100 mL of toluene were combined with the
5 oil and the mixture evaporated again at 40-70°C, 20 torr, causing the oil to solidify to form the title salt. The structure of the product was confirmed by ¹H and ¹⁹F NMR.

10

Example 7

Preparation of Dilithio perfluoro(methyl-bis-trifluoromethylsulfonimidoethyl)amine
CF₃N(CF₂CF₂SO₂N⁻SO₂CF₃)₂ · 2Li⁺

15

A 500 mL flask equipped with an addition funnel and stir bar was charged with 10 g of CF₃SO₂NH₂, 45 mL of dry THF and 45 mL of dry triethylamine, then cooled to 0°C under nitrogen. 15 g of CF₃N(CF₂CF₂SO₂F)₂

20

(prepared by electrochemical fluorination of CH₃N(CH₂CH₂SO₂F)₂) was then added slowly to the stirred solution. After complete addition, the reaction was allowed to warm to room temperature and stirred for an additional 92 hours. Next, the mixture was evaporated

25

to a brown oil under reduced pressure, which was then dissolved in 300 mL of aqueous 1M LiOH. The water, triethylamine and other volatile components were then evaporated under reduced pressure and the remaining

30

solid washed with 100 mL of diethyl ether. The ether solution was filtered, then evaporated under reduced pressure to yield a brown oil, which was dissolved in a small amount (quantity just sufficient to dissolve the oil) of diethyl ether, then added to 100 mL of methylene chloride. This solution was again filtered

35

and evaporated to yield 3.5 g of a yellow solid corresponding to the title salt. The structure was confirmed by ¹H and ¹⁹F NMR.

Example 8

Preparation of Lithium bis-
5 perfluoro(dimethylaminoethyl)sulfonimide
 $\text{Li}^+ \text{N}(\text{SO}_2\text{C}_2\text{F}_4\text{N}(\text{CF}_3)_2)_2^-$

A 500 mL Fisher-Porter pressure bottle,
equipped with a magnetic stir bar and pressure head,
10 was charged with 30 g of $(\text{CF}_3)_2\text{NC}_2\text{F}_4\text{SO}_2\text{NH}_2$ (prepared by
treatment of $(\text{CF}_3)_2\text{NC}_2\text{F}_4\text{SO}_2\text{F}$ (prepared by electrochemical
fluorination of $(\text{CH}_3)_2\text{NC}_2\text{H}_4\text{SO}_2\text{F}$) with excess ammonia in
tetrahydrofuran), 27.4 g of triethylamine and 31.8 g of
 $(\text{CF}_3)_2\text{NC}_2\text{F}_4\text{SO}_2\text{F}$. The flask was sealed and heated to
15 100°C for 37 hrs with stirring. After cooling, the
volatile components of the reaction mixture were
evaporated under reduced pressure. The residue was
dissolved in 300 mL of methylene chloride and extracted
three times with 300 mL of water each time. The
20 methylene chloride solution was then dried over MgSO_4 ,
filtered, and the solvent removed under reduced
pressure. The residue was then distilled from excess
polyphosphoric acid over a temperature and pressure
range starting from 105°C and 3 torr and ending at 97°C
25 and 0.2 torr to yield 40.13 g of a pale yellow-orange
liquid which solidified on standing. This solid,
corresponding to $\text{HN}(\text{SO}_2\text{C}_2\text{F}_4\text{N}(\text{CF}_3)_2)_2$, was dissolved in
300 mL of methyl t-butyl ether, after which 20.6 g of
 Li_2CO_3 was added in small portions with stirring. After
30 stirring overnight, a small amount of additional Li_2CO_3
was added and heated briefly until the solution tested
neutral to pH paper premoistened with distilled water.
The mixture was cooled, filtered and the solvent
removed under reduced pressure. Further drying under
35 vacuum at 110°C yielded 33.5 g of a fine white powder
corresponding to the title salt. The structure of the
product was confirmed by FTIR, and ^{19}F and ^1H NMR.

Example 9

Synthesis of Lithium (trifluoromethylsulfonyl)
(allyloxyoctafluorobutylsulfonyl) imide

5 $(\text{CH}_2=\text{CHCH}_2\text{OC}_4\text{F}_8\text{SO}_2\text{N}(\text{Li})\text{SO}_2\text{CF}_3)$

A 50 mL flask was charged with dry, distilled triethylamine (15mL), dry acetonitrile (15 mL) and $\text{CF}_3\text{SO}_2\text{NH}_2$ (6.75g, 0.0081 mole), purged with nitrogen,
10 and cooled to about 0°C in an ice bath.
 $\text{CH}_2=\text{CHCH}_2\text{OC}_4\text{F}_8\text{SO}_2\text{F}$ (3.0g, 0.0082 mole, prepared essentially as described in Gard et al., J. Fluorine Chem., 49, 331(1990) from $\text{FC}(\text{O})(\text{CF}_2)_3\text{SO}_2\text{F}$, which was prepared as described in U.S. 4,425,199) was added
15 dropwise to the solution with stirring, and the reaction mixture was allowed to come to room temperature overnight.

The solvent from the resulting brown mixture was removed under reduced pressure, and aqueous LiOH
20 (30mL of 1M) was added. The mixture was stirred until neutralization was complete. The water was removed under reduced pressure and the resulting oil was dissolved in diethyl ether, then filtered to removed excess LiOH.

25 To the ether solution was added methylene chloride (25mL) and sulfuric acid (15mL of 2M). The organic layer was separated, neutralized with excess Li_2CO_3 , filtered and the insoluble portion washed with acetonitrile. The volatile components of the combined
30 filtrate solutions were removed under reduced pressure, and the resulting brown oil slowly crystallized to yield clear crystals of the title compound. The structure was confirmed by ^1H and ^{19}F NMR. Yield 2.73g, 71%.

35

Example 10

Preparation of Lithium (trifluoromethylsulfonyl)
(fluorosulfonyl)imide, $\text{CF}_3\text{SO}_2\text{NLiSO}_2\text{F}$.

- 5 A dry, 3-necked flask equipped with a nitrogen inlet, stir bar and dry-ice condensor was charged with dry triethylamine (15mL) and cooled in a dry ice/isopropanol bath. Sulfuryl fluoride SO_2F_2 (7.0g, 6.8mmole) was bubbled into the cooled triethylamine.
- 10 The reaction mixture was allowed to warm to -30°C , followed by dropwise addition of a solution of triethylamine (10mL), acetonitrile (10mL) and $\text{CF}_3\text{SO}_2\text{NH}_2$ (4.0g, 0.027 mole). The temperature was maintained for an hour, allowed to warm to room temperature, and then
- 15 stirred overnight.

- The solvent was removed from the brown reaction mixture under reduced pressure. The remaining brown oil was dissolved in methylene chloride (75mL) and washed with water, followed by 3M H_2SO_4 , then dried
- 20 over MgSO_4 . The solvent was removed under reduced pressure, and the crude product purified by Kugelrohr distillation (80°C at 1 torr) to yield $\text{CF}_3\text{SO}_2\text{NHSO}_2\text{F}$ as a clear oil. This imide was dissolved in methyl t-butyl ether, neutralized with excess Li_2CO_3 , filtered and the
- 25 solvent removed under reduced pressure to yield the title compound. Yield 3.5g, 55%.

Example 11

Preparation of Lithium

- 30 (difluoromethylsulfonyl) (trifluoromethylsulfonyl)imide,
(HCF_2SO_2) (CF_3SO_2)NLi

- Difluoromethylsulfonyl chloride (30g, 0.2 mole) was dissolved in diethyl ether (100 mL) and cooled to 0°C . An excess of anhydrous NH_3 was added
- 35 until the conversion to the difluoromethylsulfonamide

was complete by gas chromatography/mass spectroscopy. The ether solution was evaporated to yield a reddish oil, from which a yellow crystalline solid precipitated. The crystals were washed with diethyl
5 ether, dried and then washed with hot toluene to remove yellow, oil impurities.

A mixture of the purified difluoromethylsulfonamide (26.5g, 0.2 mole), triethylamine (60.6g, 0.6 mole) and
10 trifluoromethylsulfonyl fluoride (85% pure, 39.5g, 0.26 mole) were charged into a stainless steel pressure vessel and heated at 45°C for 24 hours. The lower phase was separated from the upper triethylamine phase, dissolved in methylene chloride, and washed with water.
15 The solvent was removed under reduced pressure to yield $(\text{HCF}_2\text{SO}_2)(\text{CF}_3\text{SO}_2)\text{N}^-(\text{Et})_3\text{NH}^+$ (38.9g, 0.11 mole, 55%)

This ammonium salt was acidified with concentrated H_2SO_4 (20 g) and distilled at 100°C and 5 torr to yield $(\text{HCF}_2\text{SO}_2)(\text{CF}_3\text{SO}_2)\text{NH}$ (9.8 g). This imide
20 (15.3 g, 0.06 mole from two preparations) was treated with $\text{LiOH}\cdot\text{H}_2\text{O}$ (2.5 g, 0.06 mole in 50 mL water). The solution was filtered and dried in a convection oven overnight at 100°C. Yield 14.1 g of $(\text{HCF}_2\text{SO}_2)(\text{CF}_2\text{SO}_2)\text{N}^-\text{Li}^+$.

25

Example 12

Preparation of bis(trifluoromethylsulfonyl)imide

30 A stainless steel pressure vessel was charged with triethylamine (328 g, 3.25 mole), sealed, and cooled to 0°C. Trifluoromethylsulfonyl fluoride (103.4 g of 85% purity, 0.68 mole) was added to the vessel via a gas inlet, followed by anhydrous ammonia (4.76 g,
35 0.28 mole). The mixture was heated at 65°C for 24

hours, cooled and the lower product phase decanted.
 The upper triethylamine phase was extracted with
 methylene chloride, then combined with the lower
 product phase and washed with water to remove by-
 5 product salts. The solvent was removed under reduced
 pressure to yield the crude salt: $(\text{CF}_3\text{SO}_2)\text{N}^-\text{NH}(\text{C}_2\text{H}_5)_3^+$.
 The salt was acidified with concentrated sulfuric acid
 (250 g), then purified by vacuum distillation to yield
 the title compound. Yield 64.6 g, b.p. 100°C at 5 torr.
 10 This imide was dissolved in water (100 mL), then
 treated with $\text{LiOH}\cdot\text{H}_2\text{O}$ (9.67 g, 0.23 mole). The solution
 was filtered, then dried to a white crystalline product
 in a convection oven at 100°C. Yield 64.9 g of lithium-
 bis-(trifluoromethylsulfonyl)imide. The structure was
 15 confirmed by ^{19}F NMR.

Example 13

Preparation of $\{(-\text{SO}_2(\text{CF}_2)_4\text{SO}_2\text{N}^-\text{SO}_2(\text{CF}_2)_3\text{SO}_2\text{N}^--)\cdot$
 20 $2(\text{C}_2\text{H}_5)_3\text{NH}^+\}_n$ Polymer

The reagents $\text{H}_2\text{NSO}_2(\text{CF}_2)_4\text{SO}_2\text{NH}_2$ (1.995 g, 5.539
 mmols; prepared by electrochemical fluorination of
 $\text{FSO}_2(\text{CH}_2)_4\text{SO}_2\text{F}$ followed by amidation), $\text{FSO}_2(\text{CF}_2)_3\text{SO}_2\text{F}$
 25 (1.751 g, 5.539 mmols; prepared by electrochemical
 fluorination of $\text{FSO}_2(\text{CH}_2)_3\text{SO}_2\text{F}$), triethylamine (3.363 g,
 33.23 mmols. dried over activated 3A molecular sieves)
 and anhydrous acetonitrile (5 mL) were combined in a
 dry 2 oz (~59.1 mL) jar equipped with a magnetic stir
 30 bar and a tight-fitting cap to prevent significant
 exposure to ambient moisture. Once all reagents were
 fully dissolved, the contents were heated in an oil
 bath at 65°C with stirring overnight. The temperature
 was then increased to 75°C for 8 hours after which the
 35 reaction solution was allowed to cool to room

temperature. Volatile components of the reaction mixture were removed under vacuum at 70°C. The remaining viscous residue was combined with a mixture of methylene chloride and water and agitated in a separatory funnel to extract by-product salts into the aqueous phase. The polymeric product was only partially soluble in the methylene chloride phase, the remainder separated out as a third viscous brown oil phase. The water fraction was separated and discarded, and the remaining two organic fractions were extracted an additional two times with water. The isolated organic fractions were concentrated under vacuum at 80°C to remove solvent and any other volatile components. A brown viscous residue was obtained weighing 2.2 grams. ¹H, ¹⁹F and FTIR spectroscopic analysis identified the product as the desired title polymer. Gel permeation chromatographic analysis in methyl ethyl ketone revealed a bimodal molecular weight (MW) distribution in which 59% of the product had a number average MW of 19,670 (P.I. = 1.09) and 41% of the product had a number average MW of 4,240 (P.I. = 1.12). End group analysis by ¹⁹F NMR indicated that the overall number average MW was 1,860 and that the polymer chains were terminated with sulfonamide and sulfonate end groups. The lower than expected MW is attributed to partial hydrolysis of the sulfonyl fluoride groups by trace amounts of water in the reaction mixture.

The method shown in the Examples uses relatively inexpensive reagents and provides high yields of imides in a single step, i.e. without the isolation of intermediate reaction products. Furthermore, the method allows unsymmetrical imides (imides having two different fluoroalkyl sulfonyl groups, or fluorosulfonyl groups) to be prepared. The method also allows the preparation of polymeric imides

from imides having a polymerizable functional group.

The imides prepared using the method of this invention may be used, for example, as electrolytes in fuel cells and batteries and as catalysts for polymerization and

5 esterification.

What is claimed is:


1. A process for preparing an imide salt
selected from bis(fluoroalkylsulfonyl)imide
5 and(fluoroalkylsulfonyl)(fluorosulfonyl)imide salts
comprising the step of contacting a
fluoroalkylsulfonamide with a sulfonyl halide selected
from the group consisting of fluoroalkylsulfonyl
halides and fluorosulfonyl halides in the presence of a
10 sufficient amount of a non-nucleophilic base at a
temperature and for a time sufficient to yield the
imide salt.

2. The process of Claim 1 wherein the
15 fluoroalkylsulfonamide is prepared by contacting a
fluoroalkylsulfonyl salt with anhydrous ammonia.

3. The process of Claim 2 wherein the anhydrous
ammonia is generated *in situ* by the reaction of an
20 ammonium halide (NH_4^+X^-) with a non-nucleophilic base.

4. The process of Claim 2 where the temperature
is from -25°C to 150°C .

25 5. The process of Claim 1 characterized by the
reaction scheme:



where each Z is a fluorine atom or a polymerizable
30 organic functional group, R_f and R'_f are fluoroalkylene
groups optionally containing catenary oxygen or
nitrogen, X is a halogen atom, and B is a non-
nucleophilic base.

6. The process of Claim 5 wherein either or both of R_f and R'_f are perfluoroalkylene groups of the formula $-C_mF_{2m}-$, where m is 1 to 12.

5 7. The process Claim 5 further comprising the step of isolating the bis(fluoroalkylsulfonyl)imide salts of the formula $(Z-R_f-SO_2)(Z-R'_f-SO_2)N^+BH^+$ by evaporating the volatile reaction product components, extracting the remaining reaction product in water and
10 methylene chloride, drying and filtering the methylene chloride phase, and removing residual solvent to yield a concentrated reaction product and acidifying by contacting the concentrated reaction product with a mineral acid to produce a bis(fluoroalkylsulfonyl)imide
15 of the formula $(Z-R_f-SO_2)(Z-R'_f-SO_2)N^+H^+$.

8. The process of Claim 1 wherein the non-nucleophilic base is an organic base selected from the group of trialkylamines, alkyl aryl amines, and
20 pyridines.

9. The process of Claim 1 further comprising contacting the bis(fluoroalkylsulfonyl)imide salt with an inorganic base having an alkali metal cation M' ,
25 volatilizing the non-nucleophilic base, and isolating the resulting bis(fluoroalkylsulfonyl)imide salt by filtration, evaporating volatile components from the residue, followed by solvent extraction of the filtrate and removal of the solvent.

30

10. A process for the preparation of bis(fluoroalkylsulfonyl)imides comprising the step of contacting at least two equivalents of a fluoroalkylsulfonyl halide with one equivalent of
35 ammonia and at least three equivalents of a non-

nucleophilic base at a temperature and for a time sufficient to yield the imide salt.

11. A process for preparing a
 5 bis(fluoroalkylsulfonyl)imide which comprises contacting a sulfonamidofluoroalkylenesulfonyl halide of the formula $X-SO_2-R_f-SO_2NH_2$ with a non-nucleophilic base at a temperature and for a time sufficient to produce a polymeric fluoroalkylsulfonimide having the
 10 repeat unit $\{SO_2-R_f-SO_2N^-\}_n$, wherein n is an integer greater than 1 and X is a halogen atom.

12. A process for preparing a polymeric fluoroalkylsulfonimide characterized by the reaction
 15 scheme:

$$H_2NSO_2-R_f-SO_2NH_2 + X-SO_2-R'_f-SO_2-X + 4B \Rightarrow \{SO_2-R_f-SO_2N^--SO_2-R'_f-N^-\}_n + 2BH + 2BHX$$
, wherein n is an integer greater than 1 and X is a halogen atom.

- 20 13. A process for preparing a polymeric fluoroalkylsulfonimide characterized by the reaction scheme:

$$X-SO_2-R_f-SO_2-X + NH_3 + 3B \Rightarrow \{SO_2-R_f-SO_2-N^-\}_n + 2BHX$$
,
 wherein n is an integer greater than 1, R_f is a
 25 fluoroalkylene group, B is a non-nucleophilic base and X is a halogen atom.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/19532

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07C303/38 C07C311/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 072 040 A (M. ARMAND) 10 December 1991 see the whole document ---	1
A	US 5 256 821 A (M. ARMAND) 26 October 1993 see the whole document ---	1
A	JOURNAL OF FLUORINE CHEMISTRY, vol. 72, no. 2, June 1995, LAUSANNE, CH, pages 203-204, XP002025806 D.D. DESMARTEAU: "Novel perfluorinated ionomers and ionenes" see the whole document ---	1

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

27 February 1997

Date of mailing of the international search report

07.03.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/US 96/19532

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF FLUORINE CHEMISTRY, vol. 52, no. 1, 1 April 1991, LAUSANNE, CH, pages 7-12, XP002025807 D.D. DESMARTEAU, ET AL.: "N-Fluoro-bis(trifluoromethanesulphonyl)- imide. An improved synthesis" see the whole document ---	1
A	BERICHTE DER DEUTSCHEN CHEMISCHEN GESELLSCHAFT, vol. 75, 1942, WEINHEIM DE, pages 532-536, XP002025808 B. HELFERICH, ET AL.: "Bis-(alkylsulfonsäure)-imide" see the whole document ---	1
A	JOURNAL OF ORGANIC CHEMISTRY, vol. 46, no. 5, 27 February 1981, WASHINGTON, DC, US, pages 1012-1014, XP002025809 I.P. BLEEKER, ET AL.: "1H and 13C CIDNP study of the radical rearrangement involved in the reaction of tert-butylsulphonyl chloride with N-hydroxysulphonamides" see page 1014, right-hand column ---	1
A	GB 1 309 621 A (BAYER) 14 March 1973 see examples 5,10,13,16,18 ---	1
A	INORGANIC AND NUCLEAR CHEMISTRY LETTERS, vol. 7, no. 2, 1971, OXFORD, GB, pages 171-175, XP000618718 H.W. ROESKY, ET AL.: "Darstellung von N-Trifluoromethansulfonyl-sulfonylfluorida mid und einge Reaktionen" see the whole document ---	1
P,X	DE 195 33 711 A (CENTRAL GLASS) 14 March 1996 see the whole document -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internz I Application No

PCT/US 96/19532

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5072040 A	10-12-91	FR 2645533 A CA 2030809 A DE 69003063 D DE 69003063 T EP 0419647 A ES 2044587 T WO 9011999 A JP 4501118 T	12-10-90 07-10-90 07-10-93 17-03-94 03-04-91 01-01-94 18-10-90 27-02-92
US 5256821 A	26-10-93	FR 2637284 A CA 2000142 A EP 0364340 A WO 9003968 A JP 3501860 T	06-04-90 05-04-90 18-04-90 19-04-90 25-04-91
GB 1309621 A	14-03-73	BE 766548 A DE 2021257 A FR 2091042 A NL 7105820 A	16-09-71 11-11-71 14-01-72 02-11-71
DE 19533711 A	14-03-96	JP 8081436 A FR 2724380 A	26-03-96 15-03-96